

References: What Therapists Should Actually Do for Suicidal Clients: Assessment, safety planning, and least intrusive intervention

Carmel, A., Templeton, E., Sorenson, S. M., & Logvinenko, E. (2018). Using the Linehan Risk Assessment and Management Protocol with a chronically suicidal patient: A case report. *Cognitive and Behavioral Practice, 25*(4), 449-459.

Goldman-Mellor S, Olfson M, Lidon-Moyano C, Schoenbaum M. Association of suicide and other mortality with emergency department presentation. *JAMA Netw Open. 2019; 2*(12):e1917571.

Interian, A., Chesin, M., Kline, A., Miller, R., St. Hill, L., Latorre, M., ... & Stanley, B. (2018). Use of the Columbia-Suicide Severity Rating Scale (C-SSRS) to classify suicidal behaviors. *Archives of suicide research, 22*(2), 278-294.

Linehan, M. M., Comtois, K. A., & Ward-Ciesielski, E. F. (2012). Assessing and managing risk with suicidal individuals. *Cognitive and Behavioral Practice, 19*(2), 218-232.

Pinals, D. A. (2019). Liability and patient suicide. *Focus, 17*(4), 349-354.

Posner, K., Brent, D., Lucas, C., Gould, M., Stanley, B., Brown, G., ... & Mann, J. (2008). *Columbia-suicide severity rating scale (C-SSRS)*. New York, NY: Columbia University Medical Center, 10.

Substance Abuse and Mental Health Services Administration (US); (2009. *Addressing Suicidal Thoughts And Behaviors in Substance Abuse Treatment* [Internet]. Rockville (MD): (Treatment Improvement Protocol (TIP) Series, No. 50.) Available from: <https://www.ncbi.nlm.nih.gov/books/NBK64022/>